


Soybean Rust Fungicide Use Guidelines for 2006**



Determining when to apply a fungicide for soybean rust is based on several factors, starting with whether or not soybean rust is present in your field. To help you through the decision-making process, If no rust is present, go to A) to determine the crop growth stage and the risk of infection to decide whether or not a fungicide is needed. If rust is present, proceed to B) below.

A) No rust is present in field


Crop growth stages



B) If rust is present in field- (less than 10% of leaves infected in lower canopy)

Spraying when rust can be easily found in the canopy may not provide satisfactory or economical control.

Crop growth stages



Need more information about fungicides?
See "Fungicide Basics" on reverse side

Fungicide Basics

Why are fungicides needed to manage soybean rust?

Asian soybean rust is caused by the fungus *Phakopsora pachyrhizi*. None of the currently available soybean varieties are resistant to the soybean rust fungus, so the only way to stop the development of the disease is to use fungicides. Although soybean rust can not overwinter in Michigan, spores can travel on storm systems from infected fields in the south to fields in the north. Once a viable spore lands on a leaf of a healthy plant and temperature, humidity, and leaf wetness conditions are favorable, the spore can germinate and then penetrate into the leaf and cause infection. As spores are produced on infected plants, these spores can move easily from infected plants to healthy plants by wind or rainsplash or being moved about by human activity or equipment. Fields can quickly become infected due to a repeating cycle of spore production and re-infection, causing considerable loss of yield.

Avoiding fungicide resistance

Resistance is an inherited change in a plant pathogen's susceptibility to a fungicide. Intensive use, overuse or misuse of certain fungicides can result in the development of resistance. Rotate fungicide with different mode of action categories for successive fungicide applications. Avoid making consecutive solo applications of a strobilurin or triazole. Use a protectant fungicide (P) alone for the first application, only if no rust is present, but risk is high. If rust is present (less than 10% incidence) and lesions are sporulating in the lower canopy, use a triazole.

No rust in field, soybeans at R1-R6 Risk HIGH- use:

1 st application	2 nd application (if needed)
strobilurin	triazole or premix
triazole	premix
chlorothalonil	triazole or premix
premix	triazole or premix

Protectant fungicides (P)

- Protect the plant against infection
- Must be applied before the spores land on the leaf
- Stop the infection process on the leaf surface
- Have no effect once the fungal mycelia enter the leaf
- Chlorothalonil- stops germination of spores
- Strobilurins- stop spore germination, spore penetration of leaf

Curative fungicides (C)

- Can stop or slow an infection that has already occurred
- Must be applied soon after initial infection
- Are not effective once the fungus begins to produce spores
- Triazoles have limited curative activity

Protectant/curative combinations (P/C)

- Combination protectant and curative fungicides- premixed.

Fungicide mobility in the plant

Fungicides differ in their mobility in the plant. Chlorothalonil, a contact fungicide, is not absorbed into the leaves. Strobilurins are absorbed by leaves and move through the leaf, but are not truly systemic. Triazoles are absorbed by leaves and can move through water-conducting tissues to the upper parts of the plant, but do not move downward in the plant. Good coverage (especially in the lower canopy) is needed with all fungicides

Trade names	class	Fung. type
Tilt, Bumper, PropriMax*	triazole	C
Domark, Folicur, Orius *	triazole	C
Laredo*	triazole	C
Quadris	strobilurin	P
Headline	strobilurin	P
Headline SBR*	strobilurin/triazole	P/C
Quilt*	strobilurin/triazole	P/C
Stratego*	strobilurin/triazole	P/C
Bravo, Echo	chlorothalonil	P

*Section 18 products- no more than two applications can be made with any single Section 18 active ingredient. No more than three collective Section 18 products can be applied in a single season. Domark- 1 application only; none after R5.

Rust in field, soybeans at R1-R6 < 10% rust in canopy

1 st application	2 nd application (if needed)
triazole	premix
premix	premix or triazole