

## ALFALFA and GRASS HAY

### Calendar of insect pests in alfalfa and grass hay in Michigan and Ohio

Pests are listed from early to late-season. Key species are highlighted in bold text.

Common name	Overwintering stage, location	April / May	June	July	August	Sept
<b>alfalfa weevil</b>	adults, in protected areas	larval defoliation		(adults present)		
clover root curculio	adults, in protected areas		egg laying	larval feeding on roots		
white grubs	larvae (grubs), underground	damage to stand from root feeding				
true armyworm	Southern USA, migrates north	caterpillars feed on grasses in mixed stands or pasture				
<b>potato leafhopper</b>	Southern USA, migrates north	avg arrival ~20 May	overlapping generations of nymphs and adults suck plant sap from alfalfa leaves and stems			
spittlebug	eggs, on residue	nymphs suck plant sap		adults suck plant sap		
plant bugs	adults or eggs, in protected areas		nymphs and adults suck plant sap from alfalfa leaves and stems			
aphids (usually pea aphid)	eggs?		nymphs and adults suck plant sap from alfalfa leaves and stems			
caterpillars (multiple species)	depends on species		caterpillars feed on leaves of legumes, grass, or both			
grasshoppers (multiple species)	egg clusters, underground			nymphs, then adults, feed on leaves		
blister beetles (multiple species)	larvae, in soil cells			adult beetles feed on alfalfa leaves and blossoms		
fall armyworm	Southern USA, migrates north				caterpillars defoliate alfalfa and mixed stands	
winter cutworm	larvae, under residue				caterpillars defoliate alfalfa late into fall; active in winter	

**Damage checklist to aid in scouting of alfalfa and grass hay in Michigan and Ohio**

<b>Plant part or timing</b> Type of damage or injury	alfalfa weevil	aphids	blister beetles	caterpillars	clover root curculio	fall armyworm	grasshoppers	plant bug	potato leafhopper	spittlebug	true armyworm	white grubs	winter cutworm
<b>Leaves</b>													
small holes in leaves	x			x		x							
tip feeding	x												
large holes			x	x		x	x						x
irregular, ragged leaf feeding				x		x	x						
skeletonized 'frosted' appearance	x												x
complete defoliation - alfalfa						x							x
complete defoliation - grasses						x					x		
generalized leaf yellowing		x						x					
yellow leaf margins (hopperburn)									x				
red leaf margins									x				
leaves cupped or crinkled		x						x	x				
leaf drop									x				
sticky leaves or sooty mold		x											
spittle masses										x			
webbed, rolled leaves				x									
<b>Roots</b>													
root hairs missing												x	
pruning of whole roots												x	
chewing scars on taproots					x								
chewed furrows on taproots					x								
girdling of the taproot					x								
<b>Stand</b>													
stand thinning or weediness					x	x						x	
stand loss					x	x						x	
<b>Other</b>													
reduced forage quality									x				
shorter stand life	x				x				x			x	
cantharidin toxin in cut hay			x										

## Biological and impact of insect pests in alfalfa and grass hay in Michigan and Ohio

*Terms used to describe the pest status of each insect*

- **Rarely a pest:** Unusual. May not even be present in the state.
- **Uncommon pest:** Maybe present but below damaging levels. An outbreak once a generation
- **Occasional pest:** Present, sometimes in high numbers. An outbreak once a decade
- **Important pest:** Present in most fields, potentially increasing to damaging levels every season. A common target of scouting, management programs, or insecticide use.
- **Sporadic pest:** Damaging levels occur after favorable weather patterns (such as drought) or mass movement from south to north during the season
- **Localized pest:** Damaging levels occur in specific locations under specific agronomic conditions, for example in no-till production or in older stands.

Pest (abbreviation)	Life cycle and Number of generations	Impact on the Crop	Conditions which favor infestation or damage	Pest Status in MI & OH
<b>alfalfa weevil</b>	Adults (and some eggs) overwinter and become active when temps reach 48°F (~ 200 degree days). Adults lay eggs in stems. There are 4 larval stages, with 80% of the feeding done by the 4 <sup>th</sup> / last instar. By mid-June, development is complete, and weevils pupate in spun cocoons on the plant or in residue. Adults feed for a few weeks, then go into summer dormancy in protected areas outside the field. They re-emerge to feed for a time in late summer and early fall.  One generation per year	<ul style="list-style-type: none"> <li>• Small larvae feed in the folded terminals, chewing small holes. Older larvae feed on leaves throughout the plant</li> <li>• From a distance, heavily skeletonized foliage appears white, like frost damage</li> <li>• Repeated or heavy damage can reduce stand life by 1-2 yrs or lead to weed infestations</li> </ul>	<ul style="list-style-type: none"> <li>• Weevil populations build over time in older stands because adults overwinter nearby. New fields can be infested quickly if they are adjacent to older stands</li> </ul>	<p>Important</p> <p>Pest status seems to be increasing. It's unclear if this is due to a reduction in bio-control levels.</p>
<b>aphids</b>  <i>usually pea aphid, a big species which may be yellow, green, or pink</i>	Assumed overwintering as eggs. Summer population is all female. Females give birth to 12-14 live young per day and do not mate to reproduce.  Multiple overlapping generations	<ul style="list-style-type: none"> <li>• All stages suck plant sap from stems and leaves</li> <li>• Heavy infestation can lead to stunting, curling of leaves, and weakening of plants</li> <li>• Huge numbers can slow regrowth after cutting</li> <li>•</li> </ul>	<ul style="list-style-type: none"> <li>• Nothing specific</li> </ul>	<p>Uncommon</p> <p>Unusual outbreak in central MICH in 2023</p>
<b>blister beetle</b>	Eggs are laid in the soil. Larvae of most species feed on grasshopper eggs and thus are 'beneficial' in some sense. Larvae overwinter and adults emerge in spring. The beetles are distinctive with a round head, narrow 'neck', and loose wings that may not cover the abdomen. Multiple species are found in fields in the region.	<ul style="list-style-type: none"> <li>• The body fluid of live and dead blister beetles has cantharidin, a chemical which blisters the mouth and digestive tract of livestock. Horses are very susceptible &amp; can die after eating hay contaminated when beetles are incorporated into bales at harvest</li> <li>• Cantharidin dose varies by blister beetle species</li> </ul>	<ul style="list-style-type: none"> <li>• Grasshopper outbreaks (thus a dry season) often precedes a bad blister beetle year in alfalfa</li> <li>• Beetles may be attracted to, and aggregate on, flowering alfalfa or weeds later in the season</li> </ul>	<p>Uncommon and Sporadic</p> <p>May be an issue during or after a dry season</p>
<b>caterpillars</b>  <i>cloverworm, loopers, earworm</i>	Many species of caterpillars are found in legume forages. Some overwinter in the region, others migrate from the south.	<ul style="list-style-type: none"> <li>• Caterpillars feed on leaves and stems; a few species roll or web leaves</li> </ul>	<ul style="list-style-type: none"> <li>• Nothing specific</li> </ul>	<p>Uncommon</p>

<b>Pest (abbreviation)</b>	<b>Life cycle and Number of generations</b>	<b>Impact on the Crop</b>	<b>Conditions which favor infestation or damage</b>	<b>Pest Status in MI &amp; OH</b>
<b>clover root curculio</b>	Adults overwinter and become active in early spring. Small larvae feed on root nodules, and larger larvae on lateral and taproot. Pupation is in soil. Adults feed for a few weeks, then go into summer dormancy. They re-emerge to feed for a time in late summer and early fall.  One generation per year	<ul style="list-style-type: none"> <li>• Feeding by small larvae on root nodules could reduce N fixation</li> <li>• Larger larvae create scars, tunnel roots, and girdle the taproot. The injury reduces nutrient flow and creates entry points for root pathogens</li> <li>• Damage accumulates each season. May reduce stand life</li> </ul>	<ul style="list-style-type: none"> <li>• Older stands, as injury accumulates</li> <li>• New seedings near older stands may be killed by beetles moving out of the older field</li> </ul>	Rare and Localized
<b>fall armyworm (FAW)</b>	FAW is a tropical species that cannot survive freezing temperatures. Adult moths migrate north, arriving mid to late season. Eggs are laid on leaves. Larvae climb plants to feed during the day. Pupation in soil.  1-3 generations, if it is warm enough in the late season. Larvae CANNOT overwinter in our area.	<ul style="list-style-type: none"> <li>• Larvae prefer grasses but will eat legumes. Feeding starts on leaf margins; all leaves and small stems can be consumed under heavy infestations, leaving non-host weeds as the only vegetation in the field</li> <li>• Mass numbers may move into a field from adjacent crops (corn, wheat), ditches, or turf</li> </ul>	<ul style="list-style-type: none"> <li>• Strong winds from the SW carry moths northward</li> <li>• Warm conditions in late summer into fall can lead to several generations</li> <li>• Grass hay or mixed stands are likely more attractive for egg laying</li> </ul>	Uncommon and Sporadic  A late-season outbreak in 2021 was the worst in ~30 years
<b>grasshoppers</b> <i>multiple species</i>	Eggs overwinter in the soil. Nymphs emerge in June. Feeding increases as they grow. Females lay groups of eggs in the undisturbed soil in late summer.  1 generation per year	<ul style="list-style-type: none"> <li>• Adults and nymphs chew on leaves; feeding has a ragged appearance</li> </ul>	<ul style="list-style-type: none"> <li>• Undisturbed pastures and forage fields are preferred egg-laying sites</li> <li>• A dry summer can lead to an outbreak the following year</li> </ul>	Uncommon in alfalfa  Sporadic in pastures. Usually after a dry season
<b>plant bugs</b> <i>e.g alfalfa plant bug, lygus bug, &amp; fleahopper</i>	Alfalfa plant bugs overwinter as eggs, while Lygus adults overwinter in residue and on field edges. Weeds and early-season crops like alfalfa are preferred hosts.  Probably one generation	<ul style="list-style-type: none"> <li>• In legume forage, adults and nymphs suck plant sap; leaves may be curled or stunted</li> <li>• In legumes grown for seed, feeding damages blossoms and seeds, reducing germination</li> </ul>	<ul style="list-style-type: none"> <li>• Nothing specific</li> </ul>	Uncommon
<b>potato leafhopper (PLH)</b>	Adults are carried into the region from the south on weather fronts in late May. Females insert eggs in stems. Nymphs hatch in 7-10 days, begin feeding immediately, and reach the winged adult stage in 2-3 weeks.  Multiple overlapping generations	<ul style="list-style-type: none"> <li>• Adults and nymphs lacerate and suck on leaves and stems, damaging cells and blocking vascular tissue</li> <li>• The classic symptom of feeding is tip yellowing or 'hopper burn' (this symptom may be red in some legumes)</li> <li>• Other symptoms include stunting and curling of leaves</li> <li>• Long term impacts = yield &amp; quality loss, shorter stand life</li> </ul>	<ul style="list-style-type: none"> <li>• New seedings are very vulnerable</li> <li>• PLH damage is worse under dry conditions, and leafhopper survival is probably better as well</li> </ul>	Sporadic and Important  problematic later in the season if populations become well-established early
<b>spittlebug</b> <i>meadow and two-lined</i>	Eggs hatch in spring. Nymphs of Meadow SB feed near the soil surface on forage plants or weeds and move higher as they grow. Two-lined SB, a species expanding north into our region, feeds on roots or stems near the soil surface for its entire juvenile stage. Nymphs of both species excrete and live in a spittle mass which protects them from predation and drying out. Adult spittlebugs lay eggs in late summer.  One generation per year	<ul style="list-style-type: none"> <li>• Adults and nymphs feed on dilute xylem sap moving from the roots into the plant. They must remove a lot of fluid to get nutrients</li> <li>• Early-season feeding by nymphs can result in plant stress, stunting, bunched top growth, and yield loss</li> <li>• Losses ranging from 10-40% are reported for first-cutting, especially if combined with alfalfa weevil damage</li> </ul>	<ul style="list-style-type: none"> <li>• Nymphs are present early in the season, so first cutting alfalfa is usually the most affected stage</li> </ul>	Uncommon

Pest (abbreviation)	Life cycle and Number of generations	Impact on the Crop	Conditions which favor infestation or damage	Pest Status in MI & OH
<b>true armyworm (TAW)</b>	Adult moths migrate into the region in early spring. Eggs are laid on grassy weeds or crops, where larvae (caterpillars) feed. Larvae pupate in the soil and adult moths emerge in a week.  2 to 3 generations per year	<ul style="list-style-type: none"> <li>• Prefer to feed on the grass portion of mixed stands or in pastures, but will feed on legumes if forced to</li> <li>• Mass numbers may move into a field from adjacent crops (corn, wheat), ditches, or turf</li> </ul>	<ul style="list-style-type: none"> <li>• Nothing specific</li> </ul>	Sporadic  Outbreaks occur in years with a heavy spring flight from the south
<b>white grubs</b>  <i>multiple species</i>	Adults (scarab beetles) emerge May-July, depending on species. Eggs are laid in the soil in the summer. Grubs feed on organic matter and roots, then move down in soil profile in late fall to overwinter. In spring, annual grub species feed for a period, then pupate. June beetle grubs have a longer life cycle and may continue to feed for several more years.	<ul style="list-style-type: none"> <li>• Larvae (grubs) in general prune roots, causing wilting, deficiencies, or plant death</li> <li>• June beetle and European chafer feed in grass hay or pasture, creating dead areas</li> <li>• Asiatic garden beetle has been found in parts of alfalfa fields with a thin stand</li> </ul>	<ul style="list-style-type: none"> <li>• Populations of many grub species are higher in fields or parts of fields with sandy soil</li> </ul>	Uncommon
<b>winter cutworm</b>  <i>The adult moth is the large yellow underwing</i>  <i>The official larval name of the caterpillar, the winter cutworm, was coined in MI</i>	Winter cutworm is a European species which was first recorded in Canada in 1979.  Moths lay eggs in the summer. Caterpillars feed on numerous hosts. The cold-tolerant larvae feed well into fall. In winter, they sometimes crawl on the snow surface on sunny winter days. Larvae resume feeding very early in spring. Pupation occurs underground in May.  One generation per year	<ul style="list-style-type: none"> <li>• During outbreaks, larvae can defoliate alfalfa stands in fall. In mixed stands, they prefer to feed on alfalfa first</li> <li>• Late-season feeding reduces stubble that traps snow (thus increasing winter injury) and depletes root reserves (reducing spring growth)</li> <li>• New alfalfa seedlings planted with an oat companion crop are attractive to moths for egg laying and may be thinned</li> </ul>	<ul style="list-style-type: none"> <li>• Nothing specific</li> </ul>	Uncommon  Michigan was the first state to document economic damage by this insect in forage crops

## Management of insect pests in alfalfa and grass hay in Michigan and Ohio

A 15-inch sweep net is a must to monitor alfalfa insects like weevil and leafhopper. A supplier for nets in the region is Great Lakes IPM in Vestaburg MI, <https://www.greatlakesipm.com/>

Pest (abbreviation)	Management options and notes	Scouting recommendation	Threshold
<b>alfalfa weevil</b>	<ul style="list-style-type: none"> <li>Biological: Multiple egg, larval, and adult parasitoids (some introduced from the weevil's native range in Europe) often provide good, free control. Numerous predators eat weevils, and a fungal pathogen kills larvae under humid conditions</li> <li>Agronomic: If alfalfa is within 10 days of harvest, early cutting is the preferred way to reduce larval numbers while keeping numerous weevil parasitoids in the system. Check regrowth for survivors.</li> </ul>	<p>A sweep net is useful to detect weevil larvae</p> <p>Starting in early May, walk a pattern in the field &amp; pick 50-100 stems into a sweep net or bucket; target older stands, since weevils overwinter locally</p>	<p>Threshold:</p> <ul style="list-style-type: none"> <li>If it is more than 2 weeks until cutting: 40% of stems with feeding</li> <li>On regrowth, after early cutting: 6-8 larvae per ft<sup>2</sup></li> </ul>
<b>aphids</b> <i>usually pea aphid</i>	<ul style="list-style-type: none"> <li>Biological: Aphids are attacked by numerous predators (ladybugs, lacewings, syrphid fly larvae) &amp; parasitoids which keep populations in check. Under humid conditions, entomopathogenic fungi wipe out aphids too</li> <li>Host plant resistance: Most alfalfa varieties have some resistance to pea aphid</li> <li>Environmental: Adequate moisture (rainfall or irrigation) reduces feeding stress and increases humidity for infection by fungal pathogens. Pea aphid populations tend to decline in mid-season when temps exceed 85°F</li> </ul>	<p>Sweep netting can detect aphids colonizing fields</p> <p>Check plant stems for aphids, count # per stem</p>	<p>Guideline for alfalfa x plant height:</p> <ul style="list-style-type: none"> <li>&lt; 10 inches: At least 50 aphids per stem</li> <li>Over 10 inches: 100 aphids per stem</li> </ul> <p>Spraying is rarely justified, as biocontrol often kicks in</p>
<b>blister beetle</b>	<ul style="list-style-type: none"> <li>Agronomic: Beetles often aggregate on blossoms, so cut alfalfa prior to bloom. Crimping forage during harvest can kill beetles, so if they are present, cut forage and give them time to escape before baling</li> <li>Agronomic: First and second cutting hay has a lower chance of beetle contamination than later cuttings</li> <li>Insecticides: Chemical control is difficult since residue must last through harvest. Furthermore, dead beetles killed by insecticide may still end up harvested into bales</li> </ul>	<p>No specific recommendation</p> <p>Walk fields prior to harvest to check for aggregations of beetles</p>	<p>No specific recommendation</p>
<b>caterpillars</b> <i>cloverworm, earworm, loopers</i>	<ul style="list-style-type: none"> <li>Biological: Many predators feed on caterpillars</li> <li>Agronomic: If alfalfa is within 10 days of harvest, early cutting is the preferred way to reduce caterpillar numbers. Check regrowth for survivors</li> </ul>	<p>No specific recommendation</p>	<p>No specific recommendation</p> <p>Use guidelines for FAW or TAW</p>
<b>fall armyworm (FAW)</b>	<ul style="list-style-type: none"> <li>Biological: Predators and parasitoids kill larvae</li> <li>Agronomic: If alfalfa or hay is within 10 days of harvest, early cutting is the preferred way to reduce larval numbers. Check regrowth for survivors</li> <li>Insecticides: Applications are most effective on small larvae (less than ¼ inch). If caterpillars are invading from an adjacent field, a limited border treatment can save money</li> <li>Pesticide resistance: Note that pyrethroids may not be very effective on FAW, since it colonizes from the south where it is sprayed in multiple crops</li> </ul>	<p>No specific recommendation.</p> <p>To detect flight into the region, use bucket pheromone traps starting in mid-July</p> <p>Fall seedings are particularly vulnerable &amp; a priority for scouting</p>	<p>Guideline for small (&lt; ¼ inch) larvae: 2-3 per square ft</p>
<b>grasshoppers</b>	<ul style="list-style-type: none"> <li>Biological: Blister beetle larvae eat eggs. Insects, birds, and mammals eat nymphs &amp; adults. Fungal pathogens kill eggs and nymphs under moist, cool conditions</li> </ul>	<p>No specific recommendation</p>	<p>Guideline for hay or pasture x plant height</p> <ul style="list-style-type: none"> <li>6 inches: 8 per square yard</li> <li>over 6 inches: 16 per square yard</li> </ul>

Pest (abbreviation)	Management options and notes	Scouting recommendation	Threshold
<b>plant bugs</b>  <i>alfalfa plant bug,</i> <i>lygus bug,</i> <i>flea hopper</i>	<ul style="list-style-type: none"> <li>• Agronomic: When alfalfa is cut, adult plant bugs may move in large numbers into neighboring fields. This can be a problem for susceptible crops like sugar beet or some vegetables which may need to be monitored</li> </ul>	No specific recommendation	None  Spraying is not recommended
<b>potato leafhopper (PLH)</b>	<ul style="list-style-type: none"> <li>• Biological - A naturally occurring fungal pathogen kills PLH under favorable conditions, usually infecting by mid-August</li> <li>• Agronomic: If alfalfa is within 10 days of harvest, early cutting is the best way to manage PLH. Many eggs and nymphs will die. Check regrowth for survivors and treat only if over threshold</li> <li>• Host plant resistance: PLH-resistant hairy varieties trap nymphs and repel adults. The level of resistance varies plant by plant but overall, resistant stands can tolerate more leafhoppers than regular alfalfa</li> <li>• Insecticides: Detailed dynamic thresholds which vary with plant height, spray cost, and hay value are available in extension bulletins or online</li> </ul>	Using a sweep net, take 5 sets of 20 sweeps. Count the total # of PLH (adults and nymphs)  Hint: Mark the net handle with inches and use it to measure the stand height	Economic threshold for alfalfa, based on #PLH in 100 sweeps: <ul style="list-style-type: none"> <li>• &lt; 3 inch = 20</li> <li>• 4-7 inch = 50</li> <li>• 8-11 inch = 100</li> <li>• &gt; 12 inch = 200</li> </ul> For resistant varieties: <ul style="list-style-type: none"> <li>• New seeding, use the regular threshold</li> <li>• Older stands, use 3x the regular threshold</li> </ul>
<b>spittlebug</b>  <i>meadow and two-lined</i>	<ul style="list-style-type: none"> <li>• Biological: Spittle masses protect nymphs from predation</li> <li>• Agronomic: Nymphs usually pupate before first cutting, so early cutting may be less of an option for control</li> </ul>	No specific recommendation	Threshold: 1 or more spittle mass per stem
<b>true armyworm (TAW)</b>	<ul style="list-style-type: none"> <li>• Biological: Predators, a tachinid parasitoid, and fungal pathogens all kill armyworm larvae</li> <li>• Agronomic: If alfalfa is within 10 days of harvest, early cutting is preferred to reduce larval numbers. Check regrowth for survivors</li> <li>• Insecticides: If caterpillars are invading a forage crop from an adjacent field, a limited border treatment can be made</li> </ul>	No specific recommendation  Feeding occurs at night or on cloudy days - check for larvae or big frass pellets on the ground	Guideline for mixed stands or pasture: 4 to 6 larvae per ft <sup>2</sup>  Note: For mixed stands, both alfalfa and grass hay must be on the label
<b>white grubs</b>	<ul style="list-style-type: none"> <li>• Biological: Natural enemies, pathogens, birds, and rodents all kill grubs.</li> <li>• Agronomic: Fields and parts of fields that are sandy tend to support higher grub populations, while numbers are low elsewhere.</li> </ul> Note: it is important to identify grubs to distinguish annual species like European chafer and Asiatic garden beetles from multiyear species of June beetles	In poor stands, use a shovel to check for grubs and root pruning  Target the sandy parts of fields	None established  There are no rescue treatments in alfalfa or hay and limited options in pasture
<b>winter cutworm</b>	<ul style="list-style-type: none"> <li>• Biological: During outbreaks, numerous insects, birds, and mammals were recorded to feed on caterpillars</li> <li>• Insecticides: If caterpillars are invading a forage crop from an adjacent field, a limited border treatment can be made</li> </ul>	No specific recommendation	None established  use guideline for FAW: 2 to 3 per square foot

## Foliar Insecticides to manage insect pests of alfalfa in Michigan and Ohio

- Insecticides are grouped by active ingredient(s), which are listed alphabetically, allowing for easy comparison of products with the same chemistry.
- Application rates are listed for pests which appear on the manufacturer label. If a column is blank, the pest is not on the label. The letters in the pest columns refer to the label use rate from column two
- **NOTE: An insecticide must be registered on both alfalfa and grass to be used on intentionally-mixed stands**

Active ingredient Trade Names	Labelled rate per acre (unless stated)	alfalfa weevil	aphids	blister beetle	caterpillars	fall armyworm	grasshoppers	plant bugs	potato leafhopper	spittlebug	true armyworm	winter cutworm	Pre harvest interval (PHI) in days	Precautions and Remarks
<b>Bacillus thuringiensis (Bt)</b> Agree WG Dipel DF / XenTari Dipel ES Javelin WG	(a) 1.0 - 2.0 lbs (a) 0.5 - 2.0 lbs (a) 1.0 - 4.0 pints (a) 0.25 - 1.5 lbs				a	a					a		0	<ul style="list-style-type: none"> <li>• Labeled for alfalfa, clover, &amp; many nongrass forage crops</li> <li>• Bts are biological insecticides that must be eaten to kill, so coverage is important. Applications ideally are made when larvae are small. Check labels for rates on specific caterpillar species, larval sizes, and infestation levels</li> <li>• Some products can be used in organic production</li> </ul>
<b>carbaryl</b> Carbaryl 4L Sevin 4F Sevin XLR Plus	(a) 0.5 - 1.0 quart (b) 1.0 quart (c) 1.0 - 1.5 quart (d) 1.5 quart	d		a	c	c		c	b		c	c	7 harvest & grazing	<ul style="list-style-type: none"> <li>• Labeled for "alfalfa, clover, birdsfoot trefoil"</li> <li>• On dense growth apply in 25-40 gal water for good coverage</li> <li>• Max 1.5 quarts per cutting</li> <li>• May temporarily bleach tender foliage</li> <li>• Bee caution: Do not apply to blooming crops or weeds</li> </ul>
<b>chlorantraniliprole</b> Coragen Prevathon Shenzi 400SC Vantacor	(a) 3.5 - 7.5 oz (b) 2.0 - 5.0 oz (a) 14.0 - 20.0 oz (b) 8.0 - 20.0 oz (a) 1.7 - 3.8 oz (b) 1.0 - 2.5 oz (a) 1.2 - 2.5 oz (b) 0.7 - 1.7 oz				a	a	b				a		0	<ul style="list-style-type: none"> <li>• Labeled for "non-grass animal feeds" including alfalfa</li> <li>• Max 1 application per cutting</li> <li>• See Prevathon label for specific adjuvants and spray timings related to grasshopper control</li> </ul>
<b>chlorantraniliprole + cyhalothrin (lambda)</b> Besiege	(a) 5.0 - 8.0 oz (b) 6.0 - 10.0 oz	b	b	b	a b	b	b	b	a	b	b	a	1 forage 7 dry hay	<ul style="list-style-type: none"> <li>• Labeled for alfalfa</li> <li>• Max 1 application per cutting</li> <li>• Pest note: Check labels for specific rates x caterpillar species</li> <li>• Spray when bees are not foraging (early morning or evening)</li> </ul>

Active ingredient Trade Names	Labelled rate per acre (unless stated)	alfalfa weevil	aphids	blister beetle	caterpillars	fall armyworm	grasshoppers	plant bugs	potato leafhopper	spittlebug	true armyworm	winter cutworm	Pre harvest interval (PHI) in days	Precautions and Remarks
<b>cyfluthrin</b> Baythroid XL Tombstone Tombstone Helios	(a) 0.8 - 1.6 oz (b) 1.6 - 2.8 oz (c) 2.0 - 2.8 oz	b			a b	b	c	b	a	a	b	a	7 grazing harvest	<ul style="list-style-type: none"> <li>Labeled for alfalfa (for mixed stands, see Table 6)</li> <li>Check labels for specific rates x caterpillar species</li> <li>Fall armyworm = control of 1<sup>st</sup> &amp; 2<sup>nd</sup> instars only, less than ¼ inch</li> <li>Helios formulation has UV protection for extended residual</li> </ul>
<b>cyhalothrin (gamma)</b> Declare  Proaxis	(a) 0.77 - 1.28 oz (b) 1.02 - 1.54 oz  (a) 1.92 - 3.20 oz (b) 2.56 - 3.84 oz	b	b	b	a b	b	b	b	a	b	b	a	1 forage 7 hay	<ul style="list-style-type: none"> <li>Labeled for alfalfa (pure stands)</li> <li>Check labels for specific rates x caterpillar species</li> <li>Spray when bees are not foraging (early morning or evening)</li> </ul>
<b>cyhalothrin (lambda)</b> Warrior II w/Zeon Tech. Grizzly Too Kendo 22.8CS Lamcap II Province II Ravage II  Grizzly Z Kendo Ravage Lambda Cyhalothrin 1EC Lambda-Cy Lambda-Cy Ag Lambda Star Lambda-T Paradigm VC Silencer Willowood Lambda-Cy1EC	(a) 0.96 - 1.60 oz (b) 1.28 - 1.92 oz  (a) 1.92 - 3.20 oz (b) 2.56 - 3.84 oz	b	b	b	a b	b	b	b	a	b	b	a	1 forage 7 hay	<ul style="list-style-type: none"> <li>Many labels specify use on alfalfa (pure stands) only</li> <li>Spray when bees are not foraging (early morning or evening)</li> <li>Fall armyworm: Some labels indicate control of 1st &amp; 2nd instars only</li> <li>See label for mandatory info on spray drift management, buffer strips, and protecting aquatic habitats</li> </ul>
<b>cypermethrin (alpha)</b> Fastac EC or CS	(a) 2.2 - 3.8 oz (b) 2.8 - 3.8 oz	a	a		a	b	b	b	a	a	b	a	3	<ul style="list-style-type: none"> <li>Labeled for alfalfa (not labeled for grasses)</li> </ul>
<b>cypermethrin (zeta)</b> Mustang Maxx	(a) 2.24 - 4.0 oz (b) 2.8 - 4.0 oz	a	a		a	b	b	b	a	a	b	a	3	<ul style="list-style-type: none"> <li>Labeled for alfalfa and “nongrass animal feeds” like clover, trefoil, lupine, etc.</li> </ul>
<b>dimethoate</b> Dimate 4E Dimethoate 400 and 4EC	(a) 0.5 - 1.0 pint		a				a	a	a				10	<ul style="list-style-type: none"> <li>Labeled for alfalfa (not labeled for grasses)</li> <li>Max one application per cutting</li> <li>Highly toxic to bees. Do not apply if bees are visiting the treated area when crop or weeds are in bloom</li> </ul>
<b>flonicamid</b> BeLeaf 50SG	(a) 2.8 oz		a					a					14	<ul style="list-style-type: none"> <li>Labeled for alfalfa</li> <li>Narrow mode of action targets aphids on contact &amp; ingestion. Aphids stop feeding, but remain on plant until they dry up</li> </ul>

Active ingredient Trade Names	Labelled rate per acre (unless stated)	alfalfa weevil	aphids	blister beetle	caterpillars	fall armyworm	grasshoppers	plant bugs	potato leafhopper	spittlebug	true armyworm	winter cutworm	Pre harvest interval (PHI) in days	Precautions and Remarks
<b>flupyradifurone</b> Sivanto HL Sivanto 200 SL Sivanto Prime	(a) 3.5 - 7.0 (a) 7.0 - 10.5 oz (a) 7.0 - 14.0 oz		a						a				7	<ul style="list-style-type: none"> <li>Labeled for alfalfa (not labeled for grasses)</li> <li>Systemic insecticide, effective on sucking pests</li> </ul>
<b>GS-omega/kappa-Hxtx-Hv1a</b> Spear-Lep	(a) 1 - 2 pts				a	a					?	?	0	<ul style="list-style-type: none"> <li>Novel mode of action which may be useful on resistant Leps. Must be applied in conjunction with a low dose of Bt (see label for details). The Bt damages the caterpillar gut allowing Spear-Lep to enter the body.</li> </ul>
<b>indoxacarb</b> Steward	(a) 4.6 - 11.3 oz (b) 6.7 - 11.3 oz	b			b		a						7	<ul style="list-style-type: none"> <li>Labeled for alfalfa</li> <li>Max 11.3 oz per cutting</li> </ul>
<b>methomyl</b> Annihilate LV Nudrin LV Lannate LV Lanveer LV Annihilate SP Corrida90WSP Lannate SP Nudrin SP	(a) 1.5 - 3.0 pts (b) 3 pts (a) 0.5 - 1.0 lb (b) 1.0 lb	b	a		a	a		a			a		7	<ul style="list-style-type: none"> <li>Labeled for alfalfa</li> <li>Do not apply to dormant/ semi-dormant alfalfa when the daily temp is &lt; 50°F</li> </ul>
<b>methoxyfenozide</b> Intrepid 2F	(a) 4 - 10 oz				a	a					a		0 grazing 3 or 7 hay depends on rate, see label	<ul style="list-style-type: none"> <li>Labeled for non-grass forages (alfalfa, clover, lupin, etc.)</li> <li>Max 1 application per cutting and 32 oz per year</li> <li>Must begin applications at first sign of feeding damage</li> </ul>
<b>permethrin</b> Perm-Up 25DF Pounce 25WP Arctic 3.2 PermaStar AG Perm-Up 3.2EC	(a) 3.2 - 12.8 oz (b) 6.4 - 12.8 oz (a) 2 - 8 oz (b) 4 - 8 oz	b	a		a	a		b	b	b		a	0 or 14 depends on rate, see label	<ul style="list-style-type: none"> <li>Labeled for alfalfa. Do not apply to mixed stands with grasses or other legumes</li> <li>Spray when bees are not foraging (early morning or evening)</li> </ul>
<b>pyrethrins</b> Evergreen EC 60-6 Pyganic EC 1.4 II Pyganic 5.0	(a) 2.0 - 12.6 oz (a) 16 - 64 oz (a) 4.5 - 15.6 oz	a	a	a	a	a	a	a	a	a	a	a	0 when sprays dry	<ul style="list-style-type: none"> <li>Plant-derived insecticides that knock down insects quickly but have short residual control. Coverage is critical</li> <li>PyGanic is OMRI listed for organic crops, but Evergreen is not</li> <li>Highly toxic to bees exposed to direct treatment. Do not apply on or drift onto blooming crops or weeds</li> </ul>
<b>sulfoxaflor</b> Transform WG	(a) 0.75- 1.0 oz (b) 1.5 - 2.75 oz		a					b					7	<ul style="list-style-type: none"> <li>Labeled for alfalfa. Moves within leaf to target sucking pests</li> </ul>

## Foliar Insecticides to manage insect pests of grass hay and pastures in Michigan and Ohio

- Insecticides are grouped by active ingredient(s), which are listed alphabetically, allowing for easy comparison of products with the same chemistry
- Application rates are listed for pests which appear on the manufacturer label. If a column is blank, the pest is not on the label. The letters in the pest columns refer to the label use rate from column two
- **NOTE: An insecticide must be registered on both alfalfa and grasses to be used on intentionally-mixed stands**

Active ingredient Trade Names	Labelled rate per acre (unless stated)	caterpillars	fall armyworm	grasshoppers	spittlebug	true armyworm	white grubs	winter cutworm	Pre harvest interval (PHI) in days	Precautions and Remarks
<b>Bacillus thuringiensis (Bt)</b> Agree WG Dipel DF Javelin WG XenTari	(a) 1.0 - 2.0 lbs (a) 0.5 - 2.0 lbs (a) 0.25 - 1.5 lbs (a) 0.5 - 2.0 lbs	a				a			0	<ul style="list-style-type: none"> <li>• Labeled for grass forage, fodder, hay</li> <li>• Bts are biological insecticides that must be eaten to kill, so coverage is important. Applications ideally are made when larvae are small. Check labels for rates on specific caterpillar species, larval sizes, and infestation levels</li> <li>• Some formulations can be used in organic production</li> </ul>
<b>carbaryl</b> Carbaryl 4L Sevin 4F Sevin XLR Plus	(a) 1.0 - 1.5 quart	a	a			a			14 grazing 14 harvest	<ul style="list-style-type: none"> <li>• Labeled for pastures and grasses grown for forage, fodder, and hay</li> <li>• Bee caution: Do not apply to blooming crops or weeds</li> </ul>
<b>chlorantraniliprole</b> Coragen Prevathon Shenzi 400SC Vantacor	(a) 3.5 - 7.5 oz (b) 2.0 - 5.0 oz (a) 14.0 - 20.0 oz (b) 8.0 - 20.0 oz (a) 1.7 - 3.8 oz (b) 1.0 - 2.5 oz (a) 1.2 - 2.5 oz (b) 0.7 - 1.7 oz	a	a	b		a			0	<ul style="list-style-type: none"> <li>• Labeled for pasture and "grass forage, fodder, and hay... that will be fed on or grazed by livestock".</li> <li>• See Prevathon label for specific adjuvants and spray timings for grasshopper control</li> </ul>
<b>chlorantraniliprole + cyhalothrin</b> Besiege	(a) 5.0 - 8.0 oz (b) 6.0 - 10.0 oz	b	b	b	b	b		a	0 grazing 7 harvest	<ul style="list-style-type: none"> <li>• Labeled for pasture and "grass grown for hay or silage"</li> </ul>

Active ingredient Trade Names	Labelled rate per acre (unless stated)	caterpillars	fall armyworm	grasshoppers	spittlebug	true armyworm	white grubs	winter cutworm	Pre harvest interval (PHI) in days	Precautions and Remarks
<b>cyfluthrin</b> Baythroid XL Tombstone Tombstone Helios	(a) 1.6 - 1.9 oz (b) 2.6 - 2.8 oz	a b	b	b	a	a		a	0 grass 7 mixed stands	<ul style="list-style-type: none"> <li>Labeled for grass, "grass for hay", "grass in mixed stands with alfalfa"</li> <li>Check labels for rate x caterpillar species</li> <li>Fall armyworm = control of 1st &amp; 2nd instars only, less than ¼ inch</li> <li>Helios formulation has UV protection for extended residual</li> </ul>
<b>cyhalothrin (gamma)</b> Declare	(a) 0.77-1.28 oz (b) 1.02 - 1.54 oz	a b	b	b	b	b		a	0 grazing & forage  7 dry hay	<ul style="list-style-type: none"> <li>Labeled for pasture and "grass grown for hay or silage"</li> </ul>
<b>cyhalothrin (lambda)</b> Warrior II w/Zeon Tech. Grizzly Too Kendo 22.8CS Lamcap II Province II Ravage II  Grizzly Z Kendo Lambda Cyhalothrin 1EC Lambda-Cy Lambda-Cy Ag Lambda Star Lambda-T ParadigmVC Ravage Silencer Willowood Lambda-Cy1EC	(a) 0.96 - 1.60 oz (b) 1.28 - 1.92 oz  (a) 1.92 - 3.20 oz (b) 2.56 - 3.84 oz	b	b	b	b	b		a	0 grazing & forage  7 dry hay	<ul style="list-style-type: none"> <li>Labeled for pasture and "grass grown for hay or silage"</li> </ul>
<b>cypermethrin (zeta)</b> Mustang Maxx	(a) 2.24 - 4.0 oz (b) 2.8 - 4.0 oz	a	b	b	a	b		a	0 hay & forage	<ul style="list-style-type: none"> <li>Labeled for pasture, grass forage, and hay</li> </ul>
<b>methoxyfenozide</b> Intrepid 2F	(a) 4 - 8 oz		a			a			0 grazing 7 hay	<ul style="list-style-type: none"> <li>Labeled for grass forage, fodder, and hay</li> <li>Max 1 application per cutting and 32 oz per year</li> <li>Must begin applications at first sign of feeding damage</li> </ul>
<b>pyrethrins</b> Evergreen EC 60-6  Pyganic EC 1.4 II  Pyganic 5.0	(a) 2.0 - 12.6 oz  (a) 16 - 64 oz  (a) 4.5 - 15.6 oz	a	a	a	a	a		a	0 when sprays dry	<ul style="list-style-type: none"> <li>Plant-derived insecticides that knock down insects quickly but have very short residual control. Coverage is critical</li> <li>PyGanic is OMRI listed for organic crops, but Evergreen is not</li> <li>Highly toxic to bees exposed to direct treatment; do not apply on or drift onto blooming crops or weeds</li> </ul>
<b>spinosad</b> Blackhawk  Tracer	(a) 1.1 - 2.2 oz  (a) 1.0 - 2.0 oz	a	a			a		a	0 forage 3 hay	<ul style="list-style-type: none"> <li>Labeled for pastures, and grass crops</li> <li>Must target egg hatch and small larvae</li> </ul>