Prairie Fen Companion Plant facts

Green bulrush Scirpus atrovirens

C=4 Sedge Flowers not readily obvious



Plant type: Sedge

Bloom period: Star-like clustered seed heads that mature from July to August, and stay on the plant through the growing season.

Distinguishing characteristics: Plants are 1-3 feet tall, with dense, star-like clustered seed heads ¹/₂-1 inch across arising from a stiff stem. Leaf blades are broad (up to ³/₄ inch wide). Like other sedges, the plant is 3-sided at the base. The stem and leaves are pale green, and this species may be confused with bottlebrush sedge when not in flower or fruit. There is a prominent green membrane, or sheath, where leaves emerge from the center stem. More photos may be found at http://www.missouriplants.com/GSR/Scirpus_ atrovirens_page.html.

Habitat quality: This species does not tolerate dense shade and becomes more common in areas that are cleared of shrubs. We found this species to be very common in the first four years after clearing buckthorn.



Where can you find this plant?



Flower photo: Jennifer Anderson @ USDA-NRCS PLANTS Database Whole plant photo: Dan Tenaglia - missouriplants.com, Copyright Karen Tenaglia

MICHIGAN STATE









Developed by: Doug Landis and Anna Fiedler, MSU Department of Entomology. Funding support: National Fish and Wildlife Foundation, Lynn and Thelma MacCready Forest and Wildlife Endowment, MSU, and Hanes Trust of the Michigan Botanical Club. Partners: The Nature Conservancy, Michigan Natural Features Inventory, The Stewardship Network, Michigan DNR Landowner Incentive Program. For more information on native plants and prairie fens, go to <u>www.nativeplants.msu.edu</u>.

Where can you find this plant?













Ohio

Wisconsin

For more information on native plants and prairie fens, go to <u>www.nativeplants.msu.edu</u>

MSU is an affirmative-action, equal-opportunity employer. Michigan State University Extension programs and materials are open to all without regard to race, color, national origin, gender, gender identity, religion, age, height, weight, disability, political beliefs, sexual orientation, marital status, family status or veteran status. Issued in furtherance of MSU Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Thomas G. Coon, Director, MSU Extension, East Lansing,MI 48824. This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by MSU Extension or bias against those not mentioned.

Key description

C=

Coefficient of conservatism. This is a value that ranges from 0 for non-native invasives to 10 for plants that would only be expected to be found in undisturbed, high quality plant communities. It is a general guideline for whether the plant would be likely to be found in an intact prairie fen (not filled in with shrubs and without invasive species). However, C values may be high for some species that are not found in prairie fen but would be part of another wetland such as a marsh. They are included here because they are a widely accepted measure of habitat quality in the Midwest (http://1.usa.gov/FQAMethod).

Flower type

Classifications here follow those of Newcomb's wildflower guide (http://amzn.com/0316604429).

Number of regular parts

The flower has this number of petals or petal-like parts that are symmetrical from the flower center (radial), with each similar to the other in shape, size, and color. There may be 3 to 7 regular parts. See image at right.

Aster

These flowers have regular parts and are symmetrical from the center, but there are more than 7. Asters have a set of disc flowers in the center of the flower and a set of ray flowers outside of the disc flowers, often called petals. They are one group (genus) within the family Asteraceae, and there are many species in this genus.

Flowers not readily obvious

The flowers of plants such as grasses, sedges, and cattail are not obvious and are often confused with the fruits (seeds) of these species. While this website does not include flower descriptions for these species, they do flower.

Irregular

The flower is not symmetrical from the center but is symmetrical down a line (bilateral). See image at right.

No flowers

A number of primitive plants, including ferns, do not flower but make spores in order to reproduce.

Parts indistinguishable

These species either have parts so small their number is difficult to determine or have no petal-like parts. This group includes goldenrods, other species with small individual flowers, and plants in the family Asteraceae that have more than 7 parts, but the parts do not form distinguishable, symmetrical ray flowers (which are often called petals).

MSU is an affirmative-action, equal-opportunity employer. Michigan State University Extension programs and materials are open to all without regard to race, color, national origin, gender, gender identity, religion, age, height, weight, disability, political beliefs, sexual orientation, marital status, family status or veteran status. Issued in furtherance of MSU Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Thomas G. Coon, Director, MSU Extension, East Lansing,MI 48824. This information is for educational purposes only. Reference to commercial products or trade names does not imply endorsement by MSU Extension or bias against those not mentioned.



