Fertilizers and nutrient management for hops

Diane Brown, Michigan State University Extension





Ammoniacal Nitrogen	4.5%
Nitric Nitrogen	0.5%
Organic Nitrogen	5.0%
(derived from urea)	3.0 %
Total Nitrogen	10%
Phosphoric Acid	5.0%
(Available, derived	
from superphosphate)	
Potash	5.0%
(Water soluble from	
Potassium sulfate)	

Soils for hop production

- Loams that are well-drained, deep, and either sandy or gravelly.
- Poorly-drained soils promote root rots and should be avoided.
- Deep soils permit the hop plant's roots, which may reach a depth of 15 feet or more, to fully develop.
- Level and gently sloping fields are preferred to rolling land because irrigation systems and trellises are easier to construct and maintain on flat terrain.


Pre-plant nutrient management for hops

- Soil test!
- Correct major issues before planting
- pH 6.0 to 6.5
- Lime- season before if necessary
- Make sure all nutrients in optimum range



Soil sampling tools


- Tools: a soil probe, spade, or trowel,
- · clean plastic pail,
- sample bags and boxes, usually available from the soil laboratory;
- a pen or marker



How to take soil samples

- Each sample should have relatively the same soil texture, topography, organic matter and cropping history
- Sample depth: 12-15 inches





How to take soil samples

- Sample in a zigzag fashion, no more than 10 A/sample
- Need 15-20 subsamples
- Combine all samples into the plastic pail and mix thoroughly.
- Fill a soil sample box with the sample, or package about a pint of soil
- Fill out an information form and send it in with the soil sample to the soil testing lab

Soil sampling



- Test at the same time of year
- Soil pH tends to be higher in spring than fall
- Extractable nutrients tend to be lower in fall after harvest
- Preplant, first couple of years, then every 3 years
- Tissue testing every 1-2 years
- Nutrient levels and pH tend to be more stable in soils with higher CEC
- CEC less than 6me/100g- K, Ca and Mg may change more rapidly

Established hops

- Don't need to sample soils every year
- Monitor changes in pH and organic matter over multiple years-
- These impact nutrient availability in the soil





What to test

- Soil pH
- Phosphorus
- Potassium
- Calcium
- Magnesium
- Zinc
- Boron
- Manganese
- Organic matter
- C.E.C.



DATE	LAB#	COU	NIY	Previou	Previous Crop		FIELD ID		SI	OIL	TEXTURE			
11/14/2013	171098	Berr	ien	Н	ops		Back	Hopyar	d M	ineral]		
SOIL NUTRIENT LEVELS			Relov	v Optimu	m (Optimum			Above Optimum					
¹ Soil pH 7.5				l build	оришиш оришиш				Aoove Optimum					
2Phosphorus	(P) 66		pm											
Potassium (F		i	opm											
3Magnesium (Mg) 93	,	pm									1		
												_		
ADDITIONAL		Optional Tests: Micronutrients (ppm) Organic Nitrate-N												
Calcium (Ca)	(meg/100 g)	% of E	changeable !	Bases Ca	в.	ronutrients Mn	(ppm) Zn		1					
(ppm)			Mg		В 1	Cu			Fe	Matter 9	ppm ppm	1		
1232	7.0	1.2	11.0	87.8			2.5	2.6	_			┙		
RECOMMENT	ATTONS.											1		
							-	illogo	Donthy	9 inoba		u .		
Limestone: NONE Tillage Depth: 8 inches														
Farget pH =	6.0						9	6 Stan	d:					
Plant Nutrie	nts:							Micronutrient: (Optional)						
Year Crop			Expected	Nitro	gen Pho	sphate	Potassium	Be	ron Ma	nganese	Zinc	Соррег		
			Yield	(lb N	(A) (lb l	P ₂ O ₉ /A)	(lb K ₂ O/A) (Ib	B/A) (lb	Mn/A)	(lb Zn/A) (lb Cu/A		
1 Hops				10	10	0	80			0.0	0.0			

_			PERTEN		enti-		us.		Date	10-bep-12		
Company Transport Transpor			HOPS TRATION HOPS HOPS N BUILD 7 88 66 14				Previous Analyses & Appl				and the same of	
100000	E SATURATION RE pressure (20 to 20%) pressure (2 to 20%) menure (2 to 20%) menure (2 to 20%) or flores (contact)	100%	76.13 15.07 2.40 2.40 2.40 4.36 4.36	PO	ORGA	· ·	Г		FOR	CONVENT	-	
:	MTROGEN HARRY	Blot Value		Antennal Pearling Man, these pa Pearling Man, these pa composes	-	to to to		URBAL PAR (AM)	0-34 III			
1	SAPATE -S	Value Found	16	BULFUR HI FOTS (MI	**			CAPUS NO	452 M	74		
-	PHOOPHARES IN PARTY LINEAR	Description of the Control of the Co	760 636 -114					j.,		II.		
	CALCIUM	Desired Value Value Floore Deficit Surpose	2362 2309 1347	NONE	1	11913	-	ties	-	Ares	400	
:	MAGNESSIAN	Segret Value Value Found Certolituspine	260 260 636	NONE	-		Г					
:	POTALISMA	Depret Value Value Franci Cartot Surplus	443 179 -013	NOT SIAFATE GO SE PH	210		POT	CAPATE O	90 pt	298		
	SODIUM	Desired Value Value Found Desired Surpose	17			***	F	***		**11		
123461	Book States	111	0.66 611 63 640 6.60	MORAN 11% MORE BULL 25% CU SOLVATE 25% 28KC BULLFATE 20%	20			MARKE BULFA		-		
	(N) None I seek of a (N) Note the set of a (N) Note the set of a (N) None of the (N) Note the set of (N) Note the set of	The second secon	e resemption of the of \$15 beautiful to the Surface of	T IS NOT A WAR.					rier .			

What about testing compost and manure?

- Saturated media test- compost used as a growth medium
- Compost analysis- compost used as fertilizer
- Maturity test for compost (Solvita test)
- Manure testing list of accredited labs:
- http://www2.mda.state.mn.us/webapp/lis/manurelabs.jsp



Why do foliar analysis?

- More reliable than soil test for judging nutrient status of the plant
- · Check for nutrient deficiency
- Petiole analysis check levels of:
- Nitrogen (N)
- Phosphorus (P)
- Potassium (K)
- Zinc (Zn)
- Boron (B)

Foliar analysis

- Sample 1st-2nd week in June
- Most recently developed leaf
- 30-60 leaf petioles
- 5-6' height





When to do petiole analysis?

- When there are symptoms of nutrient deficiency
- collect petiole samples from bines showing leaf symptoms and from bines without symptoms (healthy or normal).
- The two samples are sent and analyzed separately for comparison purposes. This will allow you to diagnose whether or not the problem is related to nutrient status.

How to take samples for petiole analysis

- Be as consistent as possible with respect to growth stage when collecting tissue samples for nutrient analysis.
- Collect samples at the same growth stage each year.
- Monitor the same areas within specific hopyards or blocks. To do this, designate and flag specific rows within a block that are revisited yearly.

Petiole sampling

- Detach each petiole from the leaf blade immediately.
- Place petioles in a paper bag
- Label each sample and keep your own record of the following: varieties sampled, block where the samples are collected, sampling date, and conditions of hopyard.
- Prior to shipping, allow samples to dry for a day in a warm, dry, well-ventilated place

Petiole testing


- A general basis for nitrate levels from petiole testing:
- 0-6,000 ppm = low
- 6,000-10,000 ppm = normal
- 10,000+ ppm = plenty
- Conversion factor- 0.1000% =1000 ppm


Petiole testing

- P: Values over 10 ppm are likely sufficient
- Soil pH can effect availability of P
- Without the return of vines or other amendments, expect a drop of 2 ppm P per year
- K: 190 ppm is the critical value for K
- Less than this you will likely want to add anywhere from 100-200 lbs K
- Without the return of vines or other amendments, expect a drop of 40 ppm K per year.

Soil testing and tissue testing

- A & L Great Lakes Lab, Inc., 3505 Conestoga Drive Fort Wayne, IN 46808. (219) 483-4759
- Michigan State University, Soil & Plant Nutrient Lab, Plant & Soil Sciences, 1066 Bogue St. Room A81 East Lansing, Mi 48824-1325, Phone: 517-355-0218, Fax: 517-355-1732 Website: http://www.spnl.msu.edu
- Kinsey Agricultural Services, Inc. 297 County Highway 357 - Charleston, Missouri 63834 Telephone (573) 683-3880 Fax (573) 683-6227 <u>www.kinseyag.com</u> (soil only)
- There are many others- these are just a few.






Why pH matters

- soil pH affects the abundance of microorganisms.
- Bacteria are generally more prevalent in alkaline soils and fungi dominate in acidic soils.
- Microbes are responsible for the cycling of nutrients.
- The most diverse and numerous populations are found in near-neutral soils.

Adjusting Soil pH

- Easiest pre-plant
- Soils will progressively acidify with normal farming practices
- Low pH- use lime
- dolomitic lime also adds Ca and Mg
- Low pH- aluminum toxicity and P deficiency



CEC- Cation exchange capacity

- indicates of the nutrient holding capability of a soil.
- the greater the clay and organic matter contents, the higher the CEC of a soil.
- CEC is calculated by adding together the amount of soil test values of potassium, calcium, magnesium, and hydrogen held on the soil particles.

CEC- Cation exchange capacity

- Loamy sands and sands usually have a CEC less than 8.
- The CEC of sandy loams frequently falls between 8 and 12.
- Loams, clay loams and clays usually have a CEC greater than 12.
- As the soil pH changes, the CEC value will also vary somewhat. The higher the CEC, the greater the capacity of the soil to hold nutrients

Soil Organic Matter

- Release of N from organic matter.
- 20 lb N / % OM / Ac / Yr

fertility

- around 100 pounds of nitrogen per acre (lb N/acre) are removed on average during hop harvest.
- typical first-year N rates are 75 lb N/ acre; in subsequent years, 100 to 150 lb N/acre.
- Low phosphorus requirements- 20-30 lb P/A
- Potassium- 80-150 lb/A

Effect of fertilizers on soil pH

- Ammonium (NH4+) or ammonium forming fertilizers (ex. urea) will cause a decrease in soil pH over time.
- Nitrate (NO3-) sources carrying a basic cation should be less acid-forming then NH4+ fertilizers.
- The presence of Ca, Mg, K, and Na in the fertilizer will slightly increase or cause no change in soil pH.
- Elemental sulfur, ammonium sulfate, and compounds such as iron can reduce the soil pH

Nutrient Sources

- Bines and leaves returned to field-use caution
- Composts/cover crops
- Animal manure
- Organic Bagged Fertilizers
- Synthetic Fertilizers

Reading a fertilizer label- what's in a bag?

- Product or brand name
- N-P-K grade %(by weight) of the three major nutrients in a fertilizer.
- Guarantees for Total Nitrogen (N), Available Phosphate (P2O5) and Soluble Potash (K2O) Example: 12-15-24 means 12% nitrogen, 15% available phosphate, and 24% soluble potash
- Net weight
- Guaranteed analysis



G	uaranteed analysis					
Element	Agricultural Fertilizers (percent)	Specialty Fertilizers (percent)				
Calcium (Ca)	1.00	1.00				
Magnesium (Mg)	1.00	0.50				
Sulfur (S)	1.00	1.00				
Boron (B)	0.125	0.02				
Chlorine	-	0.10				
Copper (Cu)	0.05 (chelate 0.10)	0.50				
Iron (Fe)	0.10	0.10				
Manganese (Mn)	1.00	1.00				
Molybdenum (Mo)	0.04	0.04				
Zinc (Zn)	0.50 (chelate 0.125)	0.50				

Fertilizer and nutrients

- Organic or conventional?
- organic- can be difficult to supply nitrogen requirements
- USDA national organic program:
- http://www.ams.usda.gov/AMSv1.0/nop
- Template for organic USDA certification:
- http://www.ams.usda.gov/AMSv1.0/getfile?d DocName=STELPRDC5091031

Fertilizer and nutrients

- USDA national organic program:
- http://www.ams.usda.gov/AMSv1.0/nop
- OMRI approved fertilizers:
- http://www.omri.org/simple-opl-search/results/fertilizer

Organic Fertilizers

- Manures, composts, worm castings
- Often high in phosphorus
- Have them tested
- Commercially prepared bulk sources available



Organic Fertilizers- what's acceptable

• Naturally occurring fertilizers or amendments

Mined or Mineral Sources	
Material	
lime - carbonate, not hydrated or burnt	
gypsum - calcium sulfate	
rock phosphate - calcium phosphate	
greensand - potassium (0-0-7)	
potassium sulfate (0-0-50)	
potassium magnesium sulfate (0-0-21)	
basalt rock powder	
granite rock powder (5-10% K ₂ O)	

Organic Fertilizers- what's acceptable

• Naturally occurring fertilizers or amendments

Animal Derived Sources				
Material	Release			
	time			
bone meal (6-12-0)	1-4 mo.			
blood meal (12-0-0)	1- 4 mo.			
fish emulsion (5-2-2) adds micronutrients	1-4 mo.			
fish meal (10-6-2)	1-4 mo.			
feather meal varies- N content 7- 12% manure - many types	4+ mo.			
3 to 5 ft³ per year				
worm castings				

Organic Fertilizers- what's acceptable

• Naturally occurring fertilizers or amendments

Plant Derived Sour	ces				
Material	Release time				
alfalfa meal (3-0.5-3)	1- 4 mo.				
soybean meal (6-1.4-2)	1-4 mo.				
cotton seed meal (6-2-2)	1-4 mo.				
kelp meal (negligible- for trace elements	4+ mo.				
Kelp powder (1-0-4)	immed. – 1 mo				
wood ash (liming action) ~1/2 the liming	value of ag lime				
composts (typ. 1.5-3.5% N, 0.5-1% P,	Very slow				
1-2% K) watch salts!					

Approximate nutrient content of manure with 0.2% 0.5% 0.5% bedding Dairy cattle without 0.5% 0.2% 0.5% bedding with Horse 0.7% 0.2% 0.7% bedding 2.8% 2.3% 1.7% with litter Poultry without 1.7% 2.4% 1.7%

Approximately 30-50% of N available in the first year Non- composted manure- 120 day pre-harvest interval http://msue.anr.msu.edu/news/fall_manure_application_tips

Want to make your own compost?

- Compost production and use- John
 Biernbaum and Andy Fogiel, Department of
 Horticulture, Michigan State University
- www.safs.msu.edu/soilecology/pdfs/Combine d%20Compost05.doc

Synthetic Fertilizers

- Nitrogen sources
- Urea- 46-0-0- converts to NH+ in 2-3 d
 - coated ureas- sulfur coated, CoRoN, Nutralene, N-Sure
- Ammonium nitrate 33-0-0
- Ammonium sulfate- 21-0-0 highly acidifying
- Calcium nitrate 16-0-0
- Potassium nitrate 13-0-44 low salt index

Synthetic Fertilizers

Common Phosphorus sources

- Triple superphosphate 0-46-0
- Diammonium phosphate 18-46-0
- Monoammonium phosphate 11-48-0

Synthetic Fertilizers

Common potassium sources

- potassium chloride 0-0-60 or 0-0-62 not recommended
- Potassium sulfate 0-0-50
- Potassium magnesium sulfate 0-0-22 Mg-11.2%, S-22.7%
- Potassium nitrate 13-0-44
- ! Excessive potassium can lead to Mg deficiency

Hop Requirements

VARIES SLIGHTLY BY VARIETY

- 3% Nitrogen
- 2% Potassium
- 0.50% Phosphorus
- · Other important nutrients

 - Zinc



Hops Nitrogen requirements

- •60 to 150 lbs of actual N per acre
- •Apply in late May to mid June
- •Base rate of application on yields
- •Also consider soil type
 - Levels of organic matter-
 - 20 lb N / % OM / Ac / Yr

hop N requirement- (N from manure + returned bines,+ cover crops) = fertilizer N to apply

Nitrogen and OM

- Low organic matter (OM) soils (1 to 2%) a rate of 150 to 200 lb /acre of N should be applied.
- If soil OM levels are between 2 and 5% than a rate of 100 to 150 lbs of N per acre should applied.
- For organic matter levels over 5% than 80 to 100 lbs of N per acre should be applied to the crop.
- First year hops (establishment year) should only receive 75 lbs N per acre.

First Year Hop Requirements

PRODUCE 1750 LBS DM/acre

- 3.0% Nitrogen = 55 Lbs
- 2.0% Potassium = 35 Lbs
- 0.50% Phosphorus = 9 lbs
- Check Zinc and Boron levels in soil



Hop Requirements

PRODUCE 5000 LBS DM/acre

- 3.0% Nitrogen = 150 Lbs
- 2.0% Potassium = 100 Lbs
- 0.50% Phosphorus = 25 lbs

CONES 1/3 to 1/2 of DM/acre

- 3.0% Nitrogen = 75 Lbs
- 2.0% Potassium = 50 Lbs
- 0.50% Phosphorus = 12.5 lbs



Yields?

You Should Know Cone Yields

1000 lbs dry cones per acre

30 to 50% of total weight

2000 to 3000 lbs total

60 to 90 lbs of N removed



Phosphorus

- Phosphorus (0 to 80 lbs/acre)
- Will depend on Al levels in soil and pH
- Will depend on soil test levels

Watch excessive P levels in soils-ZN deficiency



Potassium

- Potassium (0 to 160 lbs/acre)
- Will depend on soil type
- Will depend on yield
- Also depends on soil levels



For potash

 Category
 Low Medium
 Optimum
 High
 V. High

 K (ppm)
 0-50
 51-100
 101-130
 131-160
 >160

 K to apply
 120-150
 80-120
 60-80
 0
 0

Boron

- Boron deficiency in hops
- Symptoms
 - Delayed emergence of shoots
 - Small, distorted, chlorotic leaves
 - Shortened internodes
 - Lots of buds on the crown at ground level



Photos: Compendium of hop diseases

Boron

- Boron deficiency in hops-Based on soil test results
- < 1.5 ppm apply 1.0-1.5 lb/A
- > 1.5 ppm no need to apply
- Can be toxic if applied in excess!

Zinc

- Deficiency symptoms-
- Chlorotic leaves
- Long shoots with very small, cupped, with deeply cut lobes
- weak lateral and bine growth,
- Acid, sandy soils low in organic matter neutral to alkaline soils or high in P

Photos: Compendium of hop diseases

Zinc

- Foliar application of zinc sulfate (0.15-0.18%).
- or If Zinc is low add 2 to 4 lbs per acre
- Will need to put through irrigation or blend with other fertilizer

Information sources:

Heather Darby, University of Vermont, Building a Hop Industry In New England, powerpoint presentation, August, 2013 $\,$

Fertility Guidelines for Hops in the Northeast- Dr. Heather Darby, University of Vermont Extension Agronomist

Organic Farming Principles and Practices, John A. Biernbaum,

Department of Horticulture, Michigan State University,

Using composts in the home garden, Colorado Master Gardener Note#243 http://www.ext.colostate.edu/mg/gardennotes/243.html

Organic fertilizers, Colorado Master Gardener Note #234 http://www.ext.colostate.edu/mg/gardennotes/234.html

Michigan State University Soil and Plant Nutrient Laboratory

http://www.spnl.msu.edu/

