

## Cherry Fungicides for 2005

James E. Nugent, District Horticulturist, MSU Extension  
Dr. George Sundin, Dept of Plant Pathology, MSU

	Bloom	Petal fall	Shuck split	Covers	Pre-Harvest	Post-Harvest
<b>Bravo</b>						
<b>SI's Alone</b>						
<b>SI's with Captan</b>						
<b>Pristine</b>						
<b>Flint</b>						
<b>Syllit</b>						
<b>Syllit+Sulfur</b>						
<b>Sulfur + Other</b>						
<b>Rovral</b>						
<b>Vanguard<sup>1</sup></b>						
<b>Ziram<sup>2</sup></b>						
<b>Captan + Other<sup>3</sup></b>						
<b>Coppers<sup>4</sup></b>						
<b>Topsin M + Captan<sup>5</sup></b>						

### Keys

	Good at this time
	Labeled, but avoid due to resistance at this time
	Labeled, but not recommended at this time
	Not labeled at this time

- 1 Vanguard labeled for tarts only; brown rot only
- 2 Ziram alone is weak
- 3 Captan alone is weak at labeled rate; but is a good tank mix partner with SI's, Syllit, Ziram, or TopsinM
- 4 Copper recommended for organic tarts from PF through post-harvest. Not recommended for sweets after foliage develops
- 5 TopsinM - Use limited by resistance developed to Benlate